

- [1] Altman D, McShane L, Sauerbrei W, Taube S. Reporting recommendations for tumor marker prognostic studies (REMARK): explanation and elaboration. PLoS Med [Internet] 2012 [accessed 04/04/2013];9(5):Available from: <http://www.plosmedicine.org/article/info%3Adoi%2F10.1371%2Fjournal.pmed.1001216>.
- [2] Collins G. Opening up multivariable prediction models: consensus-based guidelines for transparent reporting [Internet]. In: BMJ Group Blogs. London: BMJ, 2011 [accessed 04/04/2013]. Available from: <http://blogs.bmj.com/bmj/2011/08/03/gary-collins-opening-up-multivariable-prediction-models/>
- [3] Hayden JA, van der Windt DA, Cartwright JL, Cote P, Bombardier C. Assessing bias in studies of prognostic factors. Ann Intern Med [Internet] 2013 [accessed 04/04/2013];158(4):280-6. Available from: <http://annals.org/article.aspx?articleid=1650776>.
- [4] PROBAST steering committee. Prediction risk of bias assessment tool (PROBAST) [Internet]. Kleijnen Systematic Reviews Ltd, 2013 [accessed 04/04/2013]. Available from: <http://www.systematic-reviews.com/probast>.
- [5] Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. Ann Intern Med [Internet] 2011 [accessed 04/04/2013];155(8):529-36. Available from: <http://annals.org/article.aspx?articleid=474994>.